



DAVID M. VERNICK, MD, FACS  
Chief of Otolaryngology,  
Assistant Clinical Professor, Harvard Medical School  
Beth Israel Deaconess Medical Center  
Otology, Neurotology, Skull Base Surgery

HARSHA V. GOPAL, MD, FACS  
Assistant Clinical Professor, Harvard Medical School  
Rhinology, Sinus Disorders,  
Snoring/Sleep Apnea,  
Head and Neck Surgery

CECILIA V. TRAN, MD  
Clinical Instructor, Harvard Medical School  
Rhinology, Sinus Disorders,  
Laryngology, Head and Neck Surgery

ANN STOCKWELL, MA, CCC-A  
Director of Audiology

RACHEAL RUSH, MCD, CCC-A  
Audiologist

TRACY BLAIR, MS, CCC-A  
Audiologist

KERRY COHEN, AU.D., CCC-A  
Audiologist

## POST-OP NASAL AND SINUS SURGERY INSTRUCTIONS

1. If you have splints or packing, it will be removed at your first post-op visit (1-8 days following surgery).
2. Begin irrigation of the nose immediately upon returning home (with or without splints or packing):
  - Mix 1 tsp. of non-iodized salt (kosher or sea salt) with 1 cup of clean water. If the solution burns, add 1 pinch of baking soda to each cup of water. If your town water is unreliable, buy distilled
  - Use a bulb syringe or nasal douche bottle to wash the nose two to three times a day. Do not use a spray bottle for this, as it is not forceful enough. If you bought a SaltAire bottle you may use that.
  - If your nose bleeds, continue the irrigation, but decrease the force used. Also, point the rinse bottle away from the nasal septum (the midline wall that separates the nostrils).
  - Use the right hand to irrigate the left nostril, and the left hand to irrigate the right.
3. What to expect – some or all of the following are normal:
  - Blocked nasal passages
  - Sinus headaches
  - Small amount of nasal bleeding
  - Yellow or green mucous discharge
4. The first week following surgery you may use Afrin 12-hour spray to temporarily relieve nasal blockage. It may be used up to four times a day. If there is some minor nasal bleeding, pinch the nose and lean your head slightly forward for 10 – 15 minutes. An ice pack to the nose may also help. Afrin nasal spray also decreases bleeding
5. Diet: regular – avoid excessively hot liquids
6. Do not drive within 24 hours of surgery. Follow driving safety instructions of the prescription narcotic medication.

7. Please avoid:

- Nose blowing, picking
- Straining – If you become constipated, use a laxative.
- Strenuous exercise, including contact sports
- Crossing legs for first 24 hours
- Weight lifting
- Hot shower
- Hot liquids (coffee, etc).
- Vigorous sexual activity.
- Alcohol
- Smoking
- Plane flights or other altitude changes for 2 – 3 weeks.

8. Medications:

- **Do not take:** aspirin, ibuprofen, Aleve, Midol, Motrin or any other non-steroidal anti-inflammatory medication for 2 weeks prior to surgery and 2 weeks following surgery.
- Avoid coumadin until informed otherwise by surgeon or primary care physician
- You can use Tylenol or the narcotic medication prescribed for pain
- An antibiotic may be prescribed

9. Call MD for unusual symptoms:

- Nosebleed that doesn't stop
- Headache that is more severe than before surgery
- Fever of 101.0° F
- Changes in mental status
- Visual changes

10. Light exercise (walking, exercise bike) is acceptable. Eat well and drink plenty of fluids.

**Call Cathy to Schedule your follow-up appointment  
617-383-6846**